

TIME-VARIABLE SUMMIT CRATER RADIANCE AT MT. ETNA

Pieri, D.C., Jet Propulsion Laboratory, Pasadena, California, 91109, USA (dave@etna.jpl.nasa.gov)

The radiant energy output of summit craters at Mt. Etna (Italy) has been observed using Landsat Thematic (TM) data for the period 1986-1995. The compiled data set consists of 35 TM scenes of the Etna summit, including the Northeast crater, La Bocca Nuova, La Voragine, and the Southeast crater. A variety of systematic time-variable behavior was noted, and of particular interest was the 1991-1993 eruption, where at least 12 clear-weather TM frames were acquired.

Summit crater time-variable radiance preceding the eruption was systematic and significant. Generally, the average maximum radiance of the four summit craters increased systematically during the 800 days preceding the 1991-1993 eruption. This behavior reached a maximum in the month preceding the eruption, then rapidly receded to a low background value, once the flank eruption commenced. In addition, intercrater radiance was phased and systematic. Such systematic behavior may be useful in monitoring pre-eruption activity crescendos at Etna and elsewhere, and may be indicative of coordinated magma movement within summit conduits.

This work was carried out under contract to NASA at the Jet Propulsion Laboratory, California Institute of Technology.